



Canterbury Southern Black-backed Gull/ Karoro strategy



Canterbury Southern Black-backed Gull/ Karoro Strategy.

Mike Bell and Paula Harborne

Wildlife Management International Ltd

PO Box 607

Blenheim 7240

New Zealand

www.wmil.co.nz

This report was prepared by Wildlife Management International Limited for Environment Canterbury.

October 2018

Citation:

This report should be cited as:

Bell, M.D.; Harborne, P. 2018. Canterbury Southern Black-backed Gull/ Karoro Strategy. Unpublished Wildlife Management International Technical Report to Environment Canterbury.

All photographs in this Report are copyright © WMIL unless otherwise credited, in which case the person or organization credited is the copyright holder.

Cover image: ©Oscar Thomas, Birds Online

1. INTRODUCTION

The Southern Black-backed Gull/ Karoro is widespread and super abundant. It occurs throughout all non-forest habitats in New Zealand, from coastal areas to the high country. Following agricultural development in New Zealand, there has been a population explosion. Southern Black-backed Gull/ Karoro have adapted exceptionally well to modified environments (particularly intensive agriculture), instantly utilising fishing waste and landfills as new food sources. The current national population is estimated to be over one million birds (Biswell 2005)

With such abundance this has often seen them considered as a 'nuisance species' posing threats to other native birds, farmers (through predation of livestock and its propensity to carry disease), as well as a potential health threat to humans.

Braided rivers are a natural feature of the South Island, with 64% occurring in Canterbury. Braided rivers in New Zealand support unique communities of plants and animals, many of which are threatened with extinction (O'Donnell et al. 2016). Braided rivers are under threat from a diverse range of sources, including introduced mammalian predators, native avian predators, weed invasion, water abstraction, nitrification, dams, modified flow regimes associated with electricity generation, river protection works, gravel extraction, and human disturbance (O'Donnell et al. 2016).

Increasing Southern Black-backed Gull/ Karoro have been implicated in the declines in braided river dependent shorebird populations. There is a growing body of evidence highlighting the predatory impacts gulls are having on these shorebirds.

With this in mind Environment Canterbury has commissioned this report to develop a Canterbury Southern Black-backed Gull/ Karoro strategy. In development of this report we review the current knowledge of Southern Black-backed Gull/ Karoro biology and assess their threats to shorebirds, along with the other impacts they have on human health. We also review the current management methods being utilised in New Zealand. As part of this review we consulted widely, encouraging input from a diverse range of sectors. This consultation included a number of public meetings along with other specific meetings targeting agencies currently undertaking Southern Black-backed Gull/ Karoro control.

This report summarises the results of this review and is divided into four parts. Part one provides an overview of Southern Black-backed Gull/ Karoro biology, habits, population numbers and trends, and the gull's legal status. Part two of the report delves into the effects and impacts of Southern Black-backed Gull/ Karoro, both on the endemic threatened shorebirds, but also on human health, farming, urban infrastructure and airport safety. Part three then brings these issues back into Canterbury context. Investigating the significance of Canterbury's braided rivers, the current status and distribution of Southern Black-backed Gull/ Karoro in the region and why a regional policy is warranted. Part four of the report provides recommended options for the development of a regional strategy for Southern Black-backed Gull/ Karoro.

2. The Southern Black-backed Gull/ Karoro

2.1 Southern Black-backed Gull/ Karoro *Larus dominicanus*

New Zealand status: Unprotected native

Conservation status: Not Threatened

Other common names: Dominican Gull, Kelp Gull (Australia), Black-backed Gull

The Southern Black-backed Gull/ Karoro is one of the country's most abundant, widespread and familiar bird species (Figure 1). The instantly recognisable large adult black and white gulls occur throughout all non-forest habitats in New Zealand, from coastal areas to high country farms. They have adapted exceptionally well to modified environments, instantly utilising fishing waste and landfills as new food sources.

A native to New Zealand, the same species (referred to as Dominican, or Kelp Gulls), are found in similar latitudes of the southern hemisphere, including Australia and South America, and their sub-Antarctic offshore islands.

Following agricultural development in New Zealand, there has been a rapid population increase. The current national population is thought to be over one million birds (Biswell 2005). This increase has often seen them considered as a 'nuisance species' posing threats to other native birds, farmers (through predation of livestock and its propensity to carry disease), as well as a potential health threat to humans.



Figure 1. An adult Southern Black-backed Gull/ Karoro; one of the most widespread and recognisable bird species in New Zealand. Image © Rebecca Bowater FPSNZ www.floraandfauna.co.nz, Birds Online.

2.2 Identification

The Southern Black-backed Gull/ Karoro is New Zealand largest gull, measuring 60cm in length, weighing approximately 1kg with a wing span of 128-142cm. Adults have a white head, neck, underparts and tail, and a black back and upper wings (except for a narrow trailing white edge). They have yellow bills with a red spot at the tip of the lower mandible, pale yellow eyes with an orange outer ring and pale greenish yellow legs (Figure 1). Juveniles look very different, with dark brown plumage, black bills and legs in their first year. Their brown feathers have pale edges giving a mottled grey-brown appearance, which gradually lightens with age and seasonal wear, until they moult into adult plumage by year four (Higgins and Davis 1996, Heather & Robertson 2005). Males are slightly bigger, however there is no visual difference between the sexes.

2.3 Habitat

In the New Zealand, Southern Black-backed Gull/ Karoro can be found inhabiting a variety of environments including, estuaries, harbours, beaches, rocky shores, riverbeds, farmland, and sparsely over subalpine tussock land. They readily occur in towns and cities, utilising sports fields, roof tops, landfills and other manmade structures. They rarely venture far out at sea and usually forage within ten kilometres of the shore. Breeding usually takes place on steep headlands, sea cliffs, sand or shingle spits and many outlying islands, as well as far inland on riverbeds, near lakes and alpine tarns. They are particularly abundant around wharves, sewage outlets, seafood processing plants and inshore fishing boats, scavenging on rubbish, organic waste and fish offal.

2.4 Breeding

Southern Black-backed Gull/ Karoro breed between October and February, in large colonies or in solitary pairs (Miskelly 2013). They are monogamous, with pair bonds continued from one breeding season to the next. The number of gulls at colonies start building up in late July, with nests constructed almost immediately (mainly by the male). The nests are a bulky collection of dry grass, small sticks, seaweed, feathers and tidal debris, often built in a well-vegetated site on bare sand, rock or mud. The clutch is of 2-3 eggs (69 x 47mm, 80g), which are greyish green with dark brown blotches. Eggs are laid 2-3 days apart and are incubated by both parents for 23-27 days. Parents guard the chicks until fledging age at approximately 7-8 weeks. After fledging, chicks remain with their parents for a couple of months, and juveniles up to 6 months old are seen begging for food (Heather & Robertson 2005).

Southern Black-backed Gull/ Karoro begin to breed from four years old. Annual adult survival is high (93%), with a life expectancy of 14 years but have been known to live up to 28 years (Fordham 1985; Heather & Robertson 1996).



Figure 2. A newly hatched Southern Black-backed Gull/ Karoro chick and egg in nest. Image © Glenda Rees, Birds Online.



A Southern Black-backed Gull/ Karoro juvenile bird begging for food from an adult. It takes four years for birds to transition from the brown juvenile plumage to the black and white adult plumage. Image © Raewyn Adams, Birds Online.

2.5 Diet and Behaviour

The Southern Black-backed Gull/ Karoro prey on and scavenge a wide range of marine and terrestrial invertebrates, fish, lizards, small mammals, other birds and their eggs and chicks. They are opportunistic, their diet varying with season and surrounding environment depending on the availability of food. They have been recorded killing young lambs and poultry, as well as stealing food from other seabirds such as terns and penguins (Heather & Robertson 2005). The gulls scavenge organic waste from landfills, sewage outlets and farms, as well as fish offal from fishing boats and meat processing factories. Southern Black-backed Gull/ Karoro are able to dive briefly below the water surface to obtain food and have been seen breaking open shells for molluscs by carry it to some height and dropping them onto rocks.

Nest sites and colonies are aggressively defended by the gulls with a long series of strident persistent “ki-och” calls, as well as swooping to ward off an array of predators, including humans. They are conspicuous and generally gregarious, often roosting on rooftops in cities, attracted to food sources provided inadvertently or intentionally by people. Flocks will often follow behind inshore fishing boats or ploughs cultivating farmland. They capable of flying long distances to access food resources, being recorded travelling up to 50km from nesting areas (Fordham 1967; Horton *et al.* 1983) southern black-backed gulls will shift their diets to readily available, anthropogenic foods when possible.

2.6 Population and distribution

Southern Black-backed Gull/ Karoro were recorded in New Zealand during James Cook’s voyages in 1770 and 1773 (Biswell 2005), however population numbers were likely to have been relatively small at that time (Oliver 1955). Although there is no detailed account during the very early period of European settlement it is likely that numbers were relative small at this time. Since this time, with the development of agriculture numbers have rapidly increased and their range expanded (Higgins and Davis 1996).

Since European colonisation of New Zealand SOUTHERN BLACK-BACKED GULL/ KARORO population numbers have increased rapidly (Oliver 1955, Turbott 1967). Human activities such as the development of ports, meat-processing factories, refuse dumps and farmland, are key drivers to the Southern Black-backed Gull/ Karoro population increase. Between 1961 and 1985 the population more than doubled along the south-west of the North Island (Powlesland & Robertson 1987), some colonies increasing up to 11 times from the 1940s to 1964 (Fordham 1967). The population has increased where human influences are not as marked, such as the Waitaki Basin, where between the 1960s and 1990s southern black-backed gulls increased in density and range (Maloney 1999).

Southern Black-backed Gull/ Karoro are widely distributed in New Zealand, existing in over 76% of the map squares surveyed 1999-2004 for the *Atlas of Bird Distribution in New Zealand* (Robertson *et al.* 2007). On islands off New Zealand, Southern Black-backed Gull/ Karoro can be found breeding on the Chatham, Bounty, Antipodes, The Snares, Campbell and Auckland Islands. They have also been reported breeding as far as the Ross Dependency, Kermadec Islands and Norfolk Islands (Heather & Robertson 2005). Widespread throughout the North and South Islands, there are hundreds of colonies, many exceeding 100 pairs, some even over 1000 pairs. The species is considered as New Zealand’s most common gull (Higgins & Davies 1996), and with an estimated population of over one million, it is referred to as a “super abundant” species (Miskelly 2013).

2.7 Movements and dispersal

Southern Black-backed Gull/ Karoro are capable of long-distance flights, moving around New Zealand, as well as are thought to make trans-Tasman crossings regularly (Williams *et al.* 2006).

To study the species dispersal, a total of 3,455 Southern Black-backed Gull/ Karoro chicks were banded between 1959 and 1993 at four Canterbury localities (three being river colonies and one an island colony) (Rowe 2013). Gulls from all four colonies dispersed widely throughout New Zealand, traveling

as far as Dipton in Southland (486km) and the Wairarapa in the north (282km) (Rowe 2013). Long-distance movements by southern black-backed gulls are not uncommon and have been recorded in the past; for example, 1320 km from Makarewa to Auckland (Robertson 1972), 840 km from Maitere River mouth to Masterton (Robertson 1973), and 530 km from Muriwai Beach to the Ruamahanga River in South Wairarapa (Robertson 1964a). Rowe (2013) concluded that the direction and dispersal distances of southern black-backed gulls are variable and can occur between the North and South Islands.

Daily movements of Southern Black-backed Gull/ Karoro are less well-known, although they have been known to fly up to 50km between feeding, roosting and nesting areas (Fordham 1967; Horton *et al.* 1983). In the 1970s, the diurnal dispersal of large gulls such as great black-backed gulls (*Larus marinus*) were observed in England. Flight paths appeared to follow topographical features such as river valleys, likely because feeding sites were coincidentally associated with them (Horton *et al.* 1983). Feeding sites included landfills, agricultural land and playing fields which gulls would arrive at shortly after first light. Other gulls arrived as late as mid-morning presumably after spending time at other sites and departed in the late afternoon back towards the roost. Pre-roosts assemblies were often formed at refuse tips closer to the roost (Horton *et al.* 1983).

Similarly, Oliver (1973) distinguished a correlation between the predominant flight paths Southern Black-backed Gull/ Karoro from colonies on Rangitoto Island to the location of eight open refuse tips in the coastal margins of Auckland. All of these refuse tips were within 50km of the Rangitoto colonies so were readily accessible to southern black-backed gulls for daily foraging trips. An aerial survey in 1965 counted 12,236 Southern Black-backed Gull/ Karoro in the Wellington area, 5977 (48.8%) of which located at or near refuse tips or meat works (Fordham 1968). This creates a clumped distribution and shows southern black-backed gulls flock to major 'artificial' feeding sites during the day.

2.8 Legal status of Southern Black-backed gull

The Wildlife Act 1953 is the legislation which provides protection to New Zealand wildlife. Essentially all wildlife in New Zealand is absolutely protected unless it is specified on one of five Schedules (Schedules 1 through 5).

Southern Black-backed Gull/ Karoro is listed on Schedule 5 of the Wildlife Act 1953, which means under Section 7 (Certain wildlife not protected) they are not protected. As such Southern Black-backed Gull/ Karoro are considered an unprotected species, and it is legal for any person to kill or be in possession of Southern Black-backed Gull/ Karoro.

Although Southern Black-backed Gull/ Karoro are unprotected, other legislation covering animal ill-treatment ensure that although the killing of Southern Black-backed Gull/ Karoro is legal, it must be carried out in a humane manner. Further there is legislation which covers the use of vertebrate toxic agents which prescribe the use of any toxin used in the control of Southern Black-backed Gull/ Karoro.

2.9 Tangata Whenua

Ngāi Tahu are the Māori people of the southern islands of New Zealand – Te Waipounamu. In 1998 Ngāi Tahu settled their Treaty claims with the crown, and confirmed their place as Kaitiaki of Te Waipounamu.

Ngāi Tahu considered Southern Black-backed Gull/ Karoro as a Taonga species and it was listed on the Crown's Settlement Deed. Through this Settlement the Crown acknowledged Ngāi Tahu has a special relationship with a number of bird, plant and marine mammal species; and this includes Southern Black-backed Gull/ Karoro. The settlement further states that those responsible for the management of these species are required to consult with and have particular regard to Ngāi Tahu views about any management proposals.

Southern Black-backed Gull/ Karoro eggs and chicks were a traditional food source for Ngāi Tahu, and Mahinga Kai was used in the Crown's Settlement Deed. Mahinga Kai refers to Ngāi Tahu interests in traditional food and other natural resources and the places where those resources are obtained. Ngāi Tahu remain interested in the ability to harvest Southern Black-backed Gull/ Karoro eggs and chicks.

3. Effects and impacts of Southern Black-backed Gull/ Karoro

Southern Black-backed Gull/ Karoro abundance especially in urban environments has caused many conflicts with people, including the transmission of pathogens through contamination of water sources, hazards to aircrafts, damage to buildings from nesting material and defecation, along with causing a general nuisance. Increasingly they are also being identified as a major conservation issue, proving to be a significant predator of threatened endemic bird species.

3.1 Public health

Adapting well to modified environments, Southern Black-backed Gull/ Karoro population numbers have rapidly increased in New Zealand taking to feeding at landfills, sewage outlets, meat processing factories and farmland (Figure 3). Foraging through urban and organic waste in their hundreds creates a public health issue, as they have been shown to carry pathogenic organisms (McDiarmid 1962). Due to their opportunistic and gregarious nature, gulls are known to be important vectors for pathogens (e.g. *Salmonella*). So although they may not be the casual source, they act as efficient dispersal agents (Hatch 1996).

Southern Black backed Gull/ Karoro have been reported to carry anthropogenically derived enteric bacteria such as *Campylobacter* spp., *Salomonella* spp. and *Esherichia coli* (Fenlon 1981; Quessey and Messier 1992; Nelson *et al.*, 2008; More *et al.*, 2017) which are considered the most common causes of food-borne zoonotic. As Southern Black-backed Gull/ Karoro can travel long distances, they can effectively spread these diseases via faecal contamination of pasture and surface waters used for drinking, recreation or irrigation (Reed *et al.* 2003).

The relationship for the transmission of diseases from gulls to humans can prove difficult, however international studies suggests that gulls such as the Southern Black-backed Gull/ Karoro are important vectors (Belant 1997). Monaghan *et al.* (1985) describes the occurrence of *Salmonella* spp. in gulls (*Larus* spp.) in England, determining a substantial positive correlation between the incident of salmonellosis in the human population and the proportion of gulls tested carrying salmonellae in that urban area. In Scotland, gull faeces in public water supplies has been specified as the most credible source for disease transmission (Jones *et al.* 1978). In a second Scottish study, gull faeces contamination of water supplies was the cause of 26 occurrences of human and animal salmonellosis (Reilly *et al.* 1981). Nelson *et al.* (2008) compared populations of *Escherichia coli* among gulls (*Larus* spp.) to wastewater and landfills in the US. Using ribotyping, a genotypical bacteria source tracking method it was determined that gulls obtain faecal bacteria from wastewater and rubbish, which they may transport to recreational beaches and waters (Nelson *et al.* 2008).

Pathogens like *Salmonella* spp. have a high survival rate in aquatic environments and if water sources are contaminated by Southern Black-backed Gull/ Karoro droppings it is both an economic and public health concern (Winfiel and Groisman, 2003).



Figure 3. Southern Black-backed Gull/ Karoro scavenging as a digger spreads and buries rubbish at the Oamaru Rubbish Tip. Gulls have adapted ready to food sources such as refuse stations and from these can spread pathogens to surround rural land and urban centres. Image © Rob Morris

3.2 Farming

Southern Black-backed Gull/ Karoro have rapidly adapted to agricultural development of New Zealand. From the earliest creation of pastoral farming, gulls expanded their range to exploit this new food resource. Southern Black-backed Gull/ Karoro are now a significant element of the avifauna on farmland, and are especially abundant in areas of intensive agriculture. As such, this has often put gulls in conflict with farmers as their impacts effect farming operations.

Southern Black-backed Gull/ Karoro are often considered a pest on farmland, were some attack cast ewes and new born lambs. In areas of high gull concentrations, this can often result in significant loses of lambs in some seasons. As such most sheep farmers consider Southern Black-backed Gull/ Karoro a pest.

Salmonella Brandenburg is a disease that has caused widespread abortions and deaths in pregnant ewes since 1996 around Canterbury, Otago and Southland. In 1998, the causative organism also spread to cattle herds, leading to diarrhoea, dysentery and deaths in both adults and calves, as well as abortions in adult cattle (Clark 2001). This disease has caused major financial losses and is an important occupational hazard, with several farmers, farm workers and veterinary practitioners having contracted the disease (Clark 2000). Principal factors involved in the local spread of the disease is environmental contamination from a high number of organisms excreted by affected animals, as well as by faecal excretion of the bacteria by Southern Black-backed Gull/ Karoro.

Southern Black-backed Gull/ Karoro are known to scavenge on aborted sheep fetuses and membranes, which have high concentrations of *Salmonella* Brandenburg. The Ministry of Agriculture and Forestry funded a project to determine the likely role the gulls have, as farmers were quick to observe and hold them responsible for the local spread of the disease (Clark 2001). In September 1999, up to 25 million *Salmonella* Brandenburg organisms per gram of intestinal contents were verified in Southern Black-backed Gull/ Karoro on affected farms, as well as on non-affected neighbouring farms (Clark *et al.* 1999). Their faeces can contaminate grazing land and pollute sources of stock water (Johnston *et al.* 1979, Fenlon 1981). They have the ability to spread the disease to other farms during abortion season, when travelling long distances to feed. The infection is highly unlikely to be carried from one season to another and does not affect the Southern Black-backed Gull/ Karoro clinically.

3.3 Urban infrastructure

Southern Black-backed Gull/ Karoro are seen in many places around New Zealand as a nuisance, especially by nesting on rooftops. They can cause structural damage to buildings from nesting material and defecation, obstruct roof drainage systems with debris, defecate on nearby vehicles, and disturb the inhabitants of buildings due to noise. Roof-nesting by Southern Black-backed Gull/ Karoro typically occurs during rapid growth of colonies on natural sites in surrounding areas and is successful due to their exploitation of anthropogenic food (Belant 1997). They are observed nesting on top of buildings and on ledges mainly in coastal towns which can have an affect businesses and home owners.

Porirua business owners located between the Spicer landfill and the harbour, have complained to the City Council about southern black-backed gulls dropping waste onto their roves, such as animal parts, bones and condoms (Nicoll 2017). Southern black-backed gulls have become a financial nuisance in urban areas harassing maintenance personnel. One business has had to remove staff members from their top storey due to southern black-backed gull excrement on the rooftop causing a horrid smell and maggots to drop down inside from the ceiling (O'Neil 2013).

This issue is also observed in other parts of the world, where gulls (*Larus spp.*) are colonising on buildings and other man-made structures such as bridges, jetties, pipelines and in 1993 an oil platform in the Irish Sea. Surveys in 1976 and 1994 in Britain and Ireland have determined that gulls (*Larus spp.*) nesting on urban infrastructure have increased in terms of breeding pairs (Herring Gulls by 10% per

annum; Lesser Black-backed Gulls by 17% per annum) as well as by the number of sites colonised (Herring gulls by 5% per annum; Lesser black-backed gulls by 13% per annum) (Raven and Coulson 1997). Due to the disturbances that nesting gulls cause, the spread of gulls into urban areas is a matter of growing concern for Britain and Ireland, as well as elsewhere in Europe, North America and other parts of the world.

3.4 Airstrike risk

In many airports around the world, bird strike can pose a great risk to aircrafts. In New Zealand, Southern Black-backed Gull/ Karoro are considered one of the most hazardous bird species for aircrafts due to their comparatively large size, abundance and that they are often observed at between 1000 and 3000 feet above ground level (Robertson 1992; Avisure 2016). Airports can be in close proximity to cities, farmland, landfills and coastal environments where Southern Black-backed Gull/ Karoro are commonly found feeding. When the gulls are feeding they may cross the flight path of an aircraft and are at risk of getting sucked into the engine, causing damage to the plane.

At the Christchurch Airport, there have been 38 strikes by Southern Black-backed Gull/ Karoro, including one damaging between 1993 and 2005. An additional 57 strikes including four damaging have been by unidentified gulls and is suspected a large proportion of these is likely by Southern Black-backed Gull/ Karoro. During a two and a-half-year period in the 1990s, 74 birds were hit at Wellington Airport, as well as 362 recorded near-misses. Of these 63% were recorded Southern Black-backed Gull/ Karoro (Hutching 2006). A number of airports around New Zealand such as Wellington Airport have monitoring and control action plans in place for Southern Black-backed Gull/ Karoro at airfields.

3.5 Native birds

Southern Black-backed Gull/ Karoro have negative impacts on threatened and protected native bird species in New Zealand. They are known to harass other sea and shore birds, attempting to breed in the same place, as well as prey on their eggs and chicks. Previous studies in New Zealand have shown that Southern Black-backed Gull/ Karoro have predated the following species;

White-fronted Tern (Williams 1963)
 Australasian Gannet (Robertson 1964b)
 Black-fronted Tern (Steffens *et al.* 2012; Bell *et al.* 2018)
 New Zealand Dotterel (Willis *et al.* 2003)
 Black Stilt (Brown and Keedwell 1998)
 Black Swan (Miers and Williams 1969)
 Black-billed Gull (Thierry *et al.* 2016; Bell and Harborne, in prep)
 Banded Dotterel (BRAID website)
 Wrybill (BRAID website, Biswell 2005)
 Variable Oystercatcher (Moorhouse 2017)
 Paradise Shelduck (Biswell 2005)
 Shore Plover (Miskelly 2013)
 Fairy Tern (Miskelly 2013)

Other species have also been observed being harassed by Southern Black-backed Gull/ Karoro and it is likely that they are being predated. For example takahe chicks have been seen to cower in the remote Murchison Mountains when the gulls fly overhead (Biswell 2005).

Southern Black-backed Gull/ Karoro are a significant predator to braided river dependant birds particularly black-fronted terns and black-billed gulls, whose colonies have been known to completely collapse due to on-going Southern Black-backed Gull/ Karoro harassment. A recent study monitoring Black-fronted Tern on the Waitaki River, Canterbury, used remote cameras to film predation events (such as by mammalian predators) and concluded that Southern Black-backed Gull/ Karoro were the primary predator, responsible for 62.5% of the predation events (Schlesselmann, 2018). Predation by


Southern Black-backed Gull/ Karoro took place throughout daylight hours and although black-fronted terns were often filmed bombarding the intruder, they were unable to deter the much larger species (Schlesselmann, n.p). Similar black-fronted tern monitoring projects carried out in 2011 and 2017, filmed numerous predation events by Southern Black-backed Gull/ Karoro on the Wairau, Clarence and Acheron Rivers (Steffens 2011; Bell *et al.* 2018).

Southern Black-backed Gull/ Karoro predation can be severe, and cause total nesting failure at colonies of other braided river dependent species. In the late 1980s, Southern Black-backed Gull/ Karoro were observed predated eggs from a white-fronted tern colony wiping out 180 nests and as well as a colony of approximately 50 red-billed gulls on shell banks near Bowentown (Biswell 2005). Bell and Harborne (in prep) observed Southern Black-backed Gull/ Karoro harassing Black-billed Gull colonies on the Wairau and Clarence Rivers, grabbing chicks and predated eggs. Using time lapse photography, Southern Black-backed Gull/ Karoro were filmed taking chicks throughout daylight hours as well as during the night resulting in no chicks surviving to fledging (See case study below). A Black-billed Gull colony of 1,500 adults located on the Waimakariri River, failed due to pressure by surrounding Southern Black-backed Gull/ Karoro colonies, who were observed entering the Black-billed Gull colony repeatedly eating eggs (Thierry *et al.* 2016).

The issue of Southern Black-backed Gull/ Karoro preying on other bird species has also been observed in other parts of the world. Southern Black-backed Gull/ Karoro have been recorded in Argentina preying on the eggs and the young of cormorants, penguins, terns, Black Oystercatchers and Southern Giant Petrels (Yorio *et al.* 1998). Among the species affected, terns appear to be particularly vulnerable to gull predation, and in several studies has resulted in the loss of nest contents and the abandonment of the breeding area (by individuals as well as by the entire colony) (Yorio and Quintana 1997).



Figure 4. Southern Black-backed Gull/ Karoro preying a Black-fronted Tern nest on the Upper Clarence River. Image © Wildlife Management International Limited



Southern Black-backed Gull/ Karoro preying a chick from a Black-billed Gull colony on the Wairau River. Image © Wildlife Management International Limited

Case study: Southern black-backed gulls at Black-billed gull colonies

Using time lapse photography, Wildlife Management International Limited (WMIL) identified predators at Black-billed Gull colonies and identified Southern Black-backed Gull/ Karoro is the most significant predator (Bell and Harborne, in prep).

The Black-billed Gull is a small endemic gull, largely restricted to the braided rivers of the eastern South Island and is listed as 'Nationally Critical' by the New Zealand Threat Classification System (Robertson *et al.* 2017). To determine productivity and threats facing the species, WMIL carried out a camera study in the 2014/2015 and 2015/16 breeding seasons at the Clarence River mouth, as well as on the Wairau River in the 2016/17 and 2017/18 breeding seasons.

Video evidence showed that Southern Black-backed Gull/ Karoro appeared to target the Black-billed Gull colonies following chick hatching. Southern Black-backed Gull/ Karoro surrounded the colonies during early incubation but little predation by gulls were recorded (Mischler and Bell 2016). As chicks started to hatch, gull predation increased rapidly, suggesting that Southern Black-backed Gull/ Karoro were selectively preying Black-billed Gull chicks, not eggs (Table A).

Table A. Predators caught on camera during day-time hours at Black-billed Gull colonies on the Clarence and Wairau Rivers. Disturbance by predators are shown per month, and any disturbance that could not be identified is labelled as unknown. SBBG = Southern Black-backed Gull/ Karoro, Harrier = Swamp Harrier.

Colony	Date	Hours	Predator present (hours)			Predator present (%)		
			SBBG	Harrier	Unknown	SBBG	Harrier	Unknown
Clarence	Nov 2014	108.1	0.12	1.06	0.70	0.11%	0.98%	0.65%
	Dec 2014	132.3	42.61	0.08	0.87	32.21%	0.60%	0.66%
Clarence	Nov 2015	183.7	0.10	0.18	0.20	0.05%	0.10%	0.11%
	Dec 2015	395.9	0.10	1.05	0.87	0.03%	0.27%	0.22%
	Jan 2016	172.8	16.81	1.13	0.45	9.73%	0.65%	0.26%
Upper Wairau	Nov 2016	257.5	42.37	0.00	0.35	16.45%	0.00%	0.14%
Mid Wairau	Nov 2016	277.0	0.12	0.00	0.05	0.05%	0.00%	0.02%
	Dec 2016	327.8	10.08	0.00	0.12	3.08%	0.00%	0.04%
Lower Wairau	Nov 2017	325.8	3.60	0.05	0.38	1.10%	0.02%	0.12%

Southern Black-backed Gull/ Karoro were recorded harassing colonies throughout the day, but a significant amount of predation from gulls was also occurring at night (Figure A). Predation pressure from Southern Black-backed Gull/ Karoro caused complete breeding failure at the Clarence colony in 2014/15 and at one of the Wairau River colonies (Mid-Wairau colony) in the 2016/17 breeding season (Bell and Harborne, in prep).

Figure A. Southern Black-backed Gull/ Karoro predating Black-billed Gull colonies on the Wairau River, although most predation occurred during the day, predation also occurred at night. In the lower image predation was recorded at 2.04 AM.



4. Southern Black-backed Gull/ Karoro Control methods

With their threats to human health and welfare, and their impacts on threatened wildlife Southern Black-backed Gull/ Karoro have been the target of control operations for more than 60 years (Caithness 1968). A range of management techniques have been employed in New Zealand.

4.1 Alpha Chloralose

Alpha Chloralose is a narcotic, often used to sedate and capture wildlife (Gregory and Wilkins, 1997), however due to its strong toxic effect on avian species it is now frequently used as a control method in New Zealand. The oral toxin induces birds first into a state of dissociation, before they become motionless and sleepy, and then usually die in a sleeping position by hypothermia. Some try to make an effort to fly and recover their equilibrium, however when the drug is administered at a high dosage death will follow relatively quickly with minimal disturbance. This method of control is seemingly the most humane form of poisoning.

Alpha Chloralose poisoning has been widely used as a control method to reduce the number of nesting Southern Black-backed Gull/ Karoro in New Zealand. Prior to a poisoning operation monitoring should be carried out to determine the area, size and distribution of a gull colony, the surrounding land use, and to estimate the timing of egg-laying. Southern Black-backed Gull/ Karoro should be targeted during the peak nesting period, when the gulls are sitting on eggs and the first chicks are hatching. During this time the parent gulls' instincts keep them in close proximity to their nests, which therefore will help reduce the spread of gulls after poisoned. Performing a poisoning operation during the time of peak incubation will reduce the number of chicks needing to be humanely disposed of, as well reduce the risk of by-kill. Non-target species such as other braided river birds are less likely to enter Southern Black-backed Gull/ Karoro colonies during incubation time, as are more attached to their colony (see case study below).

To achieve the best results when using Alpha Chloralose, pre-feeding is a crucial to condition the gulls to the same type of bait that is going to be used to carry the toxin. Pre-feeding should be carried out 3-5 days immediately before the Alpha Chloralose baits are laid out, so that the gulls become accustomed to human presence and readily eat the toxic baits when they are laid out. The same carrier in a non-toxic form should be used during the pre-feeding, so the gulls are less likely to be deterred when it comes to the laying of toxic baits.

Toxic baits should be laid during calm weather, as windy conditions may blow poisoned gulls away from the colony before dying, making the recovery of carcasses laborious. As Alpha Chloralose is more effective at lower temperatures, toxic baits should ideally be laid out as close to dusk as possible. Weather can become a limitation to Alpha Chloralose operations, as a minimum 5-day good weather gap is essential. Another limitation to this control method is that it can become very labour intensive, especially during the recovery and disposal of the gull carcasses for large colonies.

Prior to carrying out an Alpha Chloralose poisoning, the appropriate approvals and Approved Handler Test Certificates are to be gained for the operation and poison handling. The adjacent land owners to the control site should be notified prior, in case any poisoned Southern Black-backed Gull/ Karoro do advance outside the target colony area. Appropriate signage must be established at all major access locations prior to the Alpha Chloralose operation and should remain until all toxic baits and poisoned gulls are retrieved.

The use of Alpha Chloralose often raises the concerns of members of the public, and almost all operations have encountered some resistance of its use. Both Alpha Chloralose and the need to control Southern Black-backed Gull/ Karoro are not well known, and as such this often creates unease from members of the public. Currently most Alpha Chloralose operations have been small scale (<1,000 birds) and localised, without the need for wide scale public notification. As such it is likely

that this has further reduced the opposition to its uses as only immediate neighbours need to be contacted.

In the first larger scale Alpha Chloralose operation planned across 30km of the Hurunui River in December 2018, wider neighbouring landowner consultation has led to a higher level of opposition, which required significantly more consultation to resolve issues and ensure the operation was approved.

Furthermore, there is growing public concern over the use of toxins in the environment. Bearing this in mind it is likely that future operations are likely to start receiving an increased amount of resistance from people due to their stanch stand against toxin use.

Case study: Waiau Toa/ Clarence River mouth Southern Black-backed/Karoro control

At the Waiau Toa/ Clarence River mouth a mixed colony of Black-billed Gull, Red-billed Gull and White-fronted Tern have been studied since the 2012 breeding seasons. Monitoring has shown that Black-billed Gull breeding success has been extremely low. Video evidence showed Southern Black-backed Gull/ Karoro continuously entering the colony taking chicks, causing complete breeding failure in the 2014/15 and 2015/16 seasons (Mischler and Bell 2016).

In order to improve breeding success Environment Canterbury contracted Wildlife Management International Limited (WMIL) to control a nearby Southern Black-backed Gull/ Karoro colony.

In preparation for the operation all nests were located and mapped. A total of 350 nests were identified. As egg laying was protracted at this colony, egg pricking was undertaken for two weeks to ensure all birds remain incubating, and no chicks hatched prior to the toxin operation.



Figure A. WMIL staff member carrying out egg pricking at the Waiau Toa/ Clarence River southern black-backed gull colony prior to the Alpha Chloralose operation, November 2017



Figure B. The distribution of Southern Black-backed Gull/ Karoro nests at the Waiau Toa/ Clarence River mouth as mapped during planning of the control operation.

Gull control was carried out using an Alpha Chloralose operation, prior to commencing this, the appropriate permissions were gained. Including landowner and neighbouring landowner consent. Warning signs were displayed at points of entry to the operational area (Figure C), and these remained in place until one month after the operation (Figure D).

Pre-feeding using non-toxic baits started on 17 November and continued until the toxic operation on 21 November (4 days pre-feeding). Each nest had 9-10 baits scattered around the rim, pre-feeding continued until the gulls were keyed in to the baits.

On 21 November non-toxic baits were replaced with toxic baits containing 10% Alpha Chloralose. These were presented to birds in the same way, with 9-10 baits at each nest. Baiting started at the top end of the colony, ensuring baits were spread throughout the colony in a timely manner. The gulls formed a feeding frenzy immediately after the first toxic baits were laid out, and all the baits were readily consumed.

As Alpha Chloralose is fast acting, poisoned Southern Black-backed Gull/ Karoro were collected two hours after the operation. With a second sweep through the area the following morning. A total of 450 gulls were collected immediately, and a further 50 the following morning (Bell, 2018). All gull carcasses were disposed of at the Blenheim Landfill in accordance with their biological waste disposal protocols.

Following the operation, a maximum count of 55 adult Southern Black-backed Gull/ Karoro were recorded within the colony area, with the operation being regarded as successful, with a resulting 90 – 95% reduction in gulls (Bell, 2018).




Figure C. Southern Black-backed Gull/ Karoro control area and locations of warning signs erected at normal points of entry to the Waiau Toa/ Clarence River mouth.

Figure D. Example of warning sign erected at normal points of entry to the Waiau Toa/ Clarence River mouth.



4.2 Shooting

Shooting has been a control method for the used to systematically reduce Southern Black-backed Gull/ Karoro numbers. The methods has some advantages over other methods as it is not critically weather dependant nor reliant on a large number of workers.

Prior to a shooting operation, monitoring needs to be carried out at the operational area to determine the activity (i.e. feeding zones, nest sites etc.), the stage of nesting, and to estimate the number of gulls present. Operations should be timed for the height of egg-laying, when the gulls are sitting on eggs or when the first chicks are hatching. During this period the parent gulls are most protective, their instincts keeping them in close proximity to the nest site, making for the most productive time of shooting. Carrying out shooting operations before large numbers of chicks are present reduces the number of chicks needing to be humanely disposed of.

Several methods can be undertaken during a shooting operation to draw gulls to the shooter, especially when nests are scattered over a large area. An approach can be to pre-feed at the site and (i.e. rabbit carcasses) to attract gulls into a known feeding zone. Using dead gulls to create curiosity is a method proved successful in previous operations, by throwing them into the air and propping them up with wire to act as decoys. Another approach is to move in and around the riverbed and re-enter the nesting site, to trigger new outbreaks of activity giving the opportunity to target gulls.

Shooting has less potential risk to non-target species, as the shooter identifies each gull shoot. Further, shooting requires no specific permissions or consents, other than operators having a valid firearms licence. Shooting has been commonly used as a follow up to toxin operations, and is likely most efficient in smaller colonies (<500 pairs).

However, shooting can cause gulls to disperse from the control area, and it appears that birds can become gun shy both over a few days of the control period, but also over successive years of control. Which reduces the efficiency of shooting.

A range of different shooting methods have been utilised-

4.2.1 Sniping

This technique is proficient for targeting gulls that are sitting/nesting at ranges of up to 250 metres. It is especially useful when targeting individual gull's occurring in separate feeding locations or post control operation of colonies. Limitations encountered include strong winds, heavy rain and that the gulls can be reluctant to land for long periods of time when other dead gulls are present on the ground.

4.2.2 Shotgun shooting

This is an efficient technique for undertaking multiple shots at gulls in flight, and has proven to be effective at reduce numbers in breeding colonies. All gulls provoked that come into range when the shooter enters the nesting area have the opportunity to be shot. Throwing a dead gull into the air and moving in and out of the colony, can attract curious gulls closer, which were not previously in shooting range. The limitation to using a shotgun is the difficulty of shooting fast flying gulls and the shooter needs to be an experienced wing shooter to ensure the operation is achieved effectivity.

4.2.3 Aerial shooting

Aerial shooting is a developing and evolving method to control Southern Black-backed Gull/ Karoro numbers, and has been used effectively in the Mackenzie Basin. It has evolved from ground shooting as gulls appeared to becoming wary following repeated ground shooting operations. Aerial shooting is effective for small scattered colonies as the operator can cover large distances in a short period of time. The use of both experienced pilots, and shooters, is paramount in ensuring a successful control operation. The location of non-target species needs to be assessed prior to operations, due to the risk of disturbance to their colonies and breeding activity.

4.3 Sterilisation of eggs

To control Southern Black-backed Gull/ Karoro sterilisation of eggs can be employed to reduce productivity. The reduction of productivity by egg sterilisation will lead to long term population declines, however this method is a long term control as it can take decades to see significant population reduction.

A considerable amount of care must be taken to keep eggs and nests intact while undertaking egg sterilisation. As this will help to prevent adult gulls to recognise their eggs are no longer viable and lay replacement clutches. Therefore reducing the need for repeated treatments per nest.

Egg sterilisation is labour intensive and requires frequent visits to the breeding colony to achieve effective control. The method is often seen by the public as more humane and preferable to other lethal control methods targeting adult birds.

There is a series of procedures can be done to treat eggs so that they do not hatch -

4.3.1 Egg puncturing

Egg puncturing is carried out by striking a heavy-gauge needle into the egg shell and membrane. The piercing can be performed by hand or done by embedding the needle onto a long stick. This will effectively sterilize it by affecting the water balance inside killing the developing embryo. Incubating gulls can sometimes recognise eggs that have been treated by puncturing, causing them to be rejected and subsequent relaying to occur (Corkhill 1970).

4.3.2 Egg shaking

Egg Shaking is carried out by vigorously shaking eggs by hand to disrupt the internal membranes. Eggs need to be shaken until internal fluids can be heard sloshing around inside. This sterilising process can be time consuming for sizeable colonies. Again some re-laying following desertion has been observed, but not to the same extent as egg puncturing (Corkhill 1970).

4.3.3 Egg pricking

Hypodermic injecting eggs with a preservative (such as formalin), is a common technique used to control gull numbers. A needle is first used to prick a small hole in the eggshell, before using a syringe to inject 5 ml of formalin (10 %) into the egg, then sealing the hole with nail polish. This both kills the developing embryo and preserves the egg. This method reduces the number of birds which recognise that their eggs have failed, so birds continue to incubate. Reducing the number of birds that abandon nests and relay.

4.3.4 Egg oiling (dipping)

Egg oiling is a sterilising treatment used to kill the embryo, while keeping the egg still intact. It involves completely coating the egg in a thin film of non-toxic mineral oil (such as paraffin oil) or vegetable oil. By dipping the egg in oil, it prevents gases from moving through the eggshell, therefore suffocating the embryo. As with egg pricking, oiling seems to cause less abandonment and re-laying.

4.4 Other possible control methods

A range of other possible control methods, or modifying existing methods could be investigated for Southern Black-backed Gull/ Karoro control.

4.4.1 DRC-1339 (Starlicide)

DRC-1339, commercially known as Starlicide, is an oral toxicant currently registered in New Zealand to control rooks (*Corvus frugilegus*) and starlings (*Sturnus vulgaris*). It has been successfully used as an avicide elsewhere in the world to control blackbirds, pigeons, crows, ravens, magpies and gulls (USDA 2001). It is used in the United States to control nesting Herring Gulls, Greater Black-backed Gulls

around coastal areas, where high gulls' populations are negatively impacting less abundant colonial waterbirds (Solman 1994).

DRC-1339 can be applied in a variety of baits including bread, walnuts, oats, maize, brown beetles and more recently macaroni (Nelson 1994). Pre-feeding is essential to achieve the most effective results, as it keys in the target species to the baits. Non-toxic baits should be in the same form and laid out in the exact fashion as the baits containing DRC-1339. The compound is a slow acting avicide and takes up to 24 hours to be completely metabolized by the target species. Once in the bloodstream it starts to weaken the birds' liver and body functions. This causes necrosis of the birds' kidneys and circulatory impairment, leading to a non-violent death from uremic poisoning and congestion of major organs.

DRC-1339 would however need to be registered to use as a control method for Southern Black-backed Gull/ Karoro in New Zealand. Therefore Alpha Chloralose would be the preferred poison to use when carrying out a poisoning operation.

4.4.2 Non-breeding season control

Most Southern Black-backed Gull/ Karoro control operations have targeted gulls at the breeding colonies during the breeding season, historically when gulls are incubating eggs before chicks start hatching. As such this leaves a small window for the control operation. Issues with weather or other operation problems can therefore impact the operation and cause delays.

As such further investigation into undertaking control at other times of the season or in foraging areas away from colonies should be investigated.

4.4.2.1 Landfill control

The control of Southern Black-backed Gull/ Karoro at landfills and other refuse sites has been employed in New Zealand, including Alpha Chloralose operations and shooting. There is a greater risk to non-target species especially in toxin operations as other species are present at landfills.

4.4.2.2 Shoulder season control

Southern Black-backed Gull/ Karoro start occupying colonies as early as August, and there would be the opportunity to undertake control operations during this pre-breeding period. This would greatly increase the time available to undertake control at breeding colonies, especially valuable when multiple colonies on a river or district are being controlled.

5. Canterbury

5.1 Braided rivers in Canterbury

Braided rivers are a natural feature of the South Island. They are rivers, which over at least part of their reach have multiple, mobile channels that flow across a gravel flood plan (Gray and Harding 2007). Braided rivers in New Zealand support unique communities of plants and animals, many of which are threatened with extinction (O'Donnell et al. 2016).

Braided rivers are under threat from a diverse range of sources, including introduced mammalian predators, native avian predators, weed invasion, water abstraction, nitrification, dams, modified flow regimes associated with electricity generation, river protection works, gravel extraction, and human disturbance (O'Donnell et al. 2016).

Braided rivers are found throughout the country, covering a land area of over 250,000ha, although this total represents only 0.9% of the country's total land area (O'Donnell et al 2016). All but 2% occurring in the South Island, with Canterbury having a staggering 64% of New Zealand's braided rivers.

5.2 The status of braided river dependent shorebirds in Canterbury

A number of endemic braided river birds have their stronghold in Canterbury, with a significant proportion of the New Zealand population breeding in Canterbury. The National population and Canterbury population for six key endemic shorebirds are listed in Table 1 below. Both national and Canterbury estimates are derived from multiple sources and expert opinion. This highlights that Canterbury holds a significant proportion of the population of many braided river dependent shorebird, especially Black Stilt, Wrybill and South Island Pied Oystercatcher.

Table 1. Conservation status (national threat ranking), national population estimate and trend, and estimated Canterbury population for endemic braided river dependent shorebird species.

Species	National threat ranking	Estimated National population	National trend	Estimated Canterbury population
South Island Pied Oystercatcher	Declining	50,000	Declining	70%
Black Stilt	Nationally Critical	100	Increasing under intensive conservation management	100%
Banded Dotterel	Nationally Vulnerable	50,000	Declining	50%
Wrybill	Nationally Vulnerable	5,000	Declining	95%
Black-billed Gull	Nationally Endangered	75,000	Declining	35%
Black-fronted tern	Nationally Endangered	5,000	Declining	60%


Using Ornithological Society of New Zealand bird atlas data Walker and Monks (2017) highlighted significant range declines in wading birds, terns and gulls in South Island braided rivers between 1969-79 and 1999-2004. There was a contraction of the breeding range on the South Island from a relatively wide distribution of inland squares to a narrow inland distribution centred on the Mackenzie Basin. They concluded that loss of habitat due to intensive land use and predation contributed to declines in braided river shorebirds, which become less widespread across inland South Island.

The Department of Conservation maintain a database of bird counts on braided rivers across Canterbury. These are a combination of river surveys carried out by the Department, Environment Canterbury and other community conservation groups. Mapping the last decade's results from these counts confirms the results of Walker and Monks (2017). Black Stilt, Wrybill, Banded Dotterel and South Island Pied Oystercatcher all have the most significant populations in the upper river reaches (figures 5-10).

5.2.1 South Island Pied Oystercatcher

South Island Pied Oystercatcher are listed as At Risk Declining (Robertson *et al* 2017), with a national population of 50,000 birds. We estimate that 70% of the population breeds on Canterbury’s rivers. South Island Pied Oystercatcher are found throughout the braided rivers of Canterbury, although the highest concentrations of birds are found in the inland river reaches (Figure 5).


Figure 5. Distribution and abundance of South Island Pied Oystercatcher recorded in braided river surveys in Canterbury 2008-2017.



5.2.2 Black Stilt

The Black Stilt is listed as Nationally Critical (Robertson *et al* 2017), with a national population of only 100 birds. The entire population is found in Canterbury, where the species is under intensive conservation management. Black Stilt are only found breeding in the Mackenzie Basin, although birds are regularly reported from the Ashley River (Figure 6).


Figure 6. Distribution and abundance of Black Stilt recorded in braided river surveys in Canterbury 2008-2017.



5.2.3 Banded Dotterel

Banded Dotterel are listed as National Vulnerable (Robertson *et al* 2017), with a national population estimated to be 50,000 birds. We estimate that 70% of the population breeds on Canterbury's rivers. Banded Dotterel are found throughout the braided rivers of Canterbury, however the highest concentrations of birds are found in the inland river reaches (Figure 7).


Figure 7. Distribution and abundance of Banded Dotterel recorded in braided river surveys in Canterbury 2008-2017.



5.2.4 Wrybill

Wrybill are listed as National Vulnerable (Robertson *et al* 2017), with a national population of approximately 5,000 birds. We estimate that 95% of the population breeds on Canterbury’s rivers. Wrybill numbers are concentrated to the inland rivers around the Mackenzie Basin, Rangitata and Rakaia Rivers. Lower numbers are found on other rivers throughout Canterbury, but at considerable lower density (Figure 8).


Figure 8. Distribution and abundance of Wrybill recorded in braided river surveys in Canterbury 2008-2017.



5.2.5 Black-billed Gull

Black-billed gull are listed as National Endangered (Robertson *et al* 2017), with a national population of 75,000 birds. We estimate that 35% of the population breeds on Canterbury’s rivers. Black-billed Gulls are found throughout the braided rivers of Canterbury, and unlike other endemic braided river dependent shorebirds there are higher numbers in the low lying coastal river reaches (Figure 9).


Figure 9. Distribution and abundance of Black-billed Gull recorded in braided river surveys in Canterbury 2008-2017.



5.2.6 Black-fronted Tern

Black-fronted Tern are listed as Nationally Endangered (Robertson *et al* 2017), with a national population estimate of 5,000 birds. We estimate that 60% of the population breeds on Canterbury's rivers. Black-fronted Tern are found throughout the braided rivers of Canterbury, although the highest concentrations of birds are found in the inland river reaches (Figure 10).

Figure 10. Distribution and abundance of Black-fronted tern recorded in braided river surveys in Canterbury 2008-2017.




5.3 Southern Black-backed Gull in Canterbury

5.3.1 Population numbers and distribution

The Southern Black-backed Gull/ Karoro population in Canterbury is extremely large, with an estimated 35,000 breeding pairs. Allowing for pre-breeding birds (all birds <4 years old) the estimated total population of Southern Black-backed Gull/ Karoro in Canterbury is 105,000 individuals.


This population is not evenly distributed across the entire region. Using data from ebird the distribution of Southern Black-backed Gull/ Karoro in Canterbury was mapped. This highlights that gull numbers are greater in low lying areas within 60km of the coast (Figure 11). Braided rivers hold a significant proportion of the breeding population, with birds dispersing to forage across adjacent farmland. Intensive agriculture in the low lying areas has allowed Southern Black-backed Gull/ Karoro numbers to explode to current levels.

Figure 11. Distribution and density of Southern Black-backed Gull/ Karoro in Canterbury mapped using data from ebird.



In November 2016 during an aerial survey of most of the braided rivers in Canterbury all Southern Black-backed Gull/ Karoro colonies were counted and mapped. Mean colony size within 50km of the coast was significantly larger (mean 185 pairs, Ttest $p=0.002981$) than colonies >50km from the coast (mean 74 pairs) (Figure 12).

Figure 12. Size of Southern Black-backed Gull/ Karoro colonies in relation to distance from the coast.



The population of Southern Black-backed Gull/ Karoro in Canterbury is certainly continuing to increase, with marked increases seen on many of the regions braided rivers. Although there is no specific data from anecdotal evidence it appears that breeding success of colonies on braided rivers is very high. This high breeding success is likely continuing to drive population increases, and enabling the Canterbury Southern Black-backed Gull/ Karoro population to continue to increase.

5.3.2 Effects of Southern Black-backed Gull/ Karoro on Canterbury's shorebirds

Southern Black-backed Gull/ Karoro are known to be impacting on shorebirds in Canterbury. Almost all braided river shorebird projects on a number of rivers throughout the region have recorded predation by Southern Black-backed Gull/ Karoro.

With the super abundant size of the population, and their widespread distribution across the region focused on braided rivers Southern Black-backed Gulls are impacting on shorebird populations on every river. Further, it is likely that these gulls are also impacting on shorebirds in other habitats in Canterbury (See case study below).

Case study: Burwood Landfill closure and impacts on Charlesworth Wetland

Burwood Landfill is located northeast of Christchurch City and covers approximately 87 hectares. From July 1984, the landfill ran as the primary refuse disposal facility for Christchurch, with approximately 230,000 tonnes of waste internment each year (AECOM 2016). The unlined waste disposal facility attracted a large number of Southern Black-backed Gull/ Karoro daily until it closed in June 2005. The Burwood Landfill's closure largely altered the distribution and foraging behaviour of Christchurch's Southern Black-backed Gull/ Karoro, which had negative impacts on a number of local river and wetland birds.

In 1985, the population of the Southern Black-backed Gull/ Karoro on the Waimakariri River was estimated at approximately 10,000 (Avisure 2016). Following the closure of the landfill the local population reduced significantly on the Waimakariri to less than half this number, with river flood events and control efforts also contributing to this decline. Numbers of Southern Black-backed Gull/ Karoro present at the landfill plummeted from 2500-3500 to 0-10, and many of these gulls relocated to nearby coastal estuaries (A. Crossland pers comm).

Christchurch's estuaries and wetlands then saw a substantial increase in predation by Southern Black-backed Gull/ Karoro on other bird species. Mallard and Scaup ducklings, Pied Stilt chicks and small native gulls' eggs and chicks were observed preyed on by Southern Black-backed Gull/ Karoro. This increase heavily suppressed the breeding success of these targeted species in areas such as the Avon-Heathcote estuary, Bexley Wetland, Brooklands Lagoon, Charlesworth Wetland and Bromley oxidation ponds.

A mixed colony of Red-billed Gull and Black-billed Gull at Charlesworth Wetland almost failed due to Southern Black-backed Gull/ Karoro predation in the 2015/16 breeding season. Southern Black-backed Gull/ Karoro from a nearby colony of approximately 20 were observed harassing the mixed colony taking chicks over several weeks. Many Pied Stilt chicks at Charlesworth Wetland were also observed being predated during this period. The on-going pressure from Southern Black-backed Gull/ Karoro resulted in Red-billed Gull and Black-billed Gull abandoning the site, and not returning in the following breeding seasons. Currently Charlesworth Wetland is home to a colony of more than 30 Southern Black-backed Gull/ Karoro which have displaced the mixed colony, with very few other wetland bird species breeding there.



Figure A. Southern Black-backed Gull/ Karoro breeding at Charlesworth Wetland.

Image and information from Christchurch City Council (A Crossland Pers. Comm.)

5.3.3 Current Southern Black-backed Gull/ Karoro control operation

The Department of Conservation, Environment Canterbury, Christchurch International Airport and a number of community conservation groups are currently undertaking or proposing Southern Black-backed Gull/ Karoro control operations on braided rivers throughout Canterbury. The majority of these operations are to protect braided river dependent birds, with Christchurch International Airport undertaking control to reduce bird strike risk to aircraft.

Current or planned (with funding already in place) control operations are occurring on the Clarence River, Waiau river, Hurunui River, Waimakiriri river, Ashburton River and Mackenzie Basin.

At present these control operations are largely ongoing independently of each other, with no regional coordination. Further, there is no forum for the sharing of results and dissemination of learnings from control operations.

5.3.4 Cost of Southern Black-backed Gull/ Karoro control operations

The current cost of Southern Black-backed Gull/ Karoro control operations was reviewed, although this review was limited by the available data on the cost of Southern Black-backed Gull/ Karoro operations. Data was only available for 16 operations; 4 Alpha Chloralose operations, 2 ground shooting operations and 10 aerial shooting operations.

From these 16 operations the average cost was \$52.8/ gull killed. However for most operations the figures provided were only operational costs, and did not include the planning (including the permissions process) phase of the operation. As such this figure does not reflect the true cost of the operation, and we estimate that the true cost is \$62/ gull culled.

5.4 Need for a Southern Black-backed Gull/ Karoro regional strategy

Southern Black-backed Gull/ Karoro numbers are super abundant in Canterbury, and all evidence indicates that they are likely to be continuing to increase and expand their range. Linked to this there is irrefutable proof that Southern Black-backed Gull/ Karoro are having a negative impact on the breeding success of braided river dependent shorebirds. This has led to a number of control operations being undertaken, of planned to be undertaken this season.

Given their widespread distribution, increasing population and the threat they pose to shorebirds, there is a clear need for a strategy to tackle this issue on a regional scale. Southern Black-backed Gull/ Karoro are increasing and impacting shorebirds on all braided rivers throughout the region, this is a regional problem not a site specific one. Throughout all discussions with stakeholders and interest groups there was overwhelming support for the need a regional plan. The present approach to Southern Black-backed Gull/ Karoro control has each operation working independently of each other, with little or no linkages.

6. Southern Black-backed Gull Strategy

Southern Black-backed Gull/ Karoro are super abundant in Canterbury, and their numbers are continuing to increase. Their current numbers, and rate of population increase are entirely derived from anthropogenic improvements in food supply. Southern Black-backed Gull/ Karoro have readily adapted to an agricultural landscape and endless food supply from poor waste management practises.

The increasing Southern Black-backed Gull/ Karoro populations are having significant negative impacts on braided river dependent shorebird populations, and are likely to be one of the key drivers in decreasing population trends in threatened shorebirds. Furthermore, burgeoning populations of Southern Black-backed Gull/ Karoro are having harmful effects on human health and public safety.

Due to these impacts, currently a range of organisations are undertaking Southern Black-backed Gull/ Karoro control operations throughout Canterbury. These programmes are happening entirely independently of each other, with no co-ordination and little communication between practitioners.

As such there is a clear need to develop a regional strategy to address the issues of a super abundant and increasing Southern Black-backed Gull/ Karoro population in Canterbury.

6.1 Southern Black-backed Gull population control

There is irrefutable proof that Southern Black-backed Gull/ Karoro are having a negative impact on braided river dependent shorebirds, and with increasing populations this is likely to exert even more pressure on shorebird populations.

Southern Black-backed Gull/ Karoro have clearly adapted well to increased food supplies from human sources, and their populations have exploded well beyond natural levels. It is their adaption to human induced change which has created an imbalance, with now super abundant Southern Black-backed Gull/ Karoro predated other shorebird species that have significantly smaller, and declining populations.

As such there is a need to undertake some form of population control. There is a number of options, which have different impacts on and costs. Here we present three control regimes for consideration.

6.1.1 Regional wide population reduction

Given their super abundance there is a clear need to manage the numbers of Southern Black-backed Gull/ Karoro in Canterbury to improve the breeding success of braided river dependent shorebirds. Current population numbers are far in excess of pre-human levels and if there was the desire to see a marked improvement in shore bird breeding success across the entire Canterbury region, significant reductions in the regional Southern Black-backed Gull/ Karoro populations are warranted.

In order to see a regional improvement in braided river dependent shorebird breeding success it is recommended to reduce the Canterbury Southern Black-backed Gull/ Karoro population to 10% of its current numbers. Over time this would mean reducing numbers to approximately 4,000 breeding pairs, and would require the culling of an estimated 101,000 birds.

This approach would require significant community engagement to ensure support. Current Southern Black-backed Gull/ Karoro control operations generate some negative feedback, and require careful management to enable them to be carried out successfully. To date all of these control operations have been small scale (single colony focused) that don't require public notification. As such opposition has been limited, and manageable. With more widespread control across entire river systems or districts this would require a higher level notification and likely generate more opposition to the control work. As such moving to a regional control strategy would require a significant increase in the level of community engagement and consultation.

A regional wide control strategy would require significant funding. A review of the costs of current control operations indicates that the current cost is \$62/per gull controlled. As such a tentative cost for regional wide Southern Black-backed Gull/ Karoro control would be in excess of \$6.3 million³.

6.1.2 Containment control

The current distribution of Southern Black-backed Gull/ Karoro in Canterbury is not even, with numbers on the inland rivers significantly fewer than on lower river stretches (see section above). In addition, braided river dependent shore bird numbers are higher in these inland rivers. Both Wallace and Monks (2017) and an analysis of DOC held river survey data have highlighted that shorebirds in the eastern South Island have retreated to higher altitude inland river reaches.

As such it is recommended that a containment control approach is adopted and that Southern Black-backed Gull/ Karoro numbers in upper river reaches are reduced to zero density, and that ongoing management is implemented to ensure that these river reaches are not re-colonised by Southern Black-backed Gull/ Karoro.

This approach would require the control of an estimated 3,000 birds and would see control operations focused on key shorebird breeding habitat. There is a number of benefits from a containment approach. Firstly this approach would likely to be more social acceptable as it would require the culling of fewer birds and is more values focused. This will mean that it is likely than any such control operation will encounter less opposition.

A containment approach would see key shorebird habitats protected and enable a more sustainable control regime. Currently Southern Black-backed Gull/ Karoro numbers are highest in areas of intensive agricultural and population centres (low lying coastal areas). With effectively an unlimited food supply gull numbers are continuing to expand, as such any control carried out in these areas may be less effective as high breeding success means culled birds are rapidly replaced. The control of gulls in upper reaches where the surround farmland is less intensively managed is likely to be more successful. These areas are currently supporting significantly fewer gulls with a more limited food supply, making it likely that gulls can be removed from these areas long term with less management.

As stated a containment approach would require the culling of an estimated 3,000 birds, but would then require ongoing management to maintain gulls at zero density. Although currently not well understood, there would almost certainly be some immigration into the containment area from the lower river reaches. Based on current estimates of gull control operations a containment approach would be estimated to cost \$186,000. However given the isolated nature of some of the upper river reaches, these costs may be an underestimate. In addition, after initial control, there would be ongoing annual costs to maintain zero densities of gulls.

6.1.3 Values based individual river control programmes

Canterbury has significant populations of braided river dependent shorebirds and currently there are a number of mammalian predator control projects operating in Canterbury aimed at improving the breeding success of these shorebirds. There is a range of projects, with a number of project leaders; including the Department of Conservation, Environment Canterbury and Community Conservation Groups.

Although a review of these projects was outside the scope of this project, most appear to have been initiated due to high shorebird values occurring on that river. As such it is recommended that Southern Black-backed Gull/ Karoro control should be carried out in conjunction with mammalian predator control on rivers deemed of high value.

As pointed out by O'Donnell et al (2016) predator control projects on braided rivers must target the entire predator guild, including native avian predators. As such any mammalian predator control project should also incorporate Southern Black-backed Gull/ Karoro control. Existing projects which do not include Southern Black-backed Gull/ Karoro control should immediately incorporate this, and

all future projects should include Southern Black-backed Gull/ Karoro control during their development, planning and fund sourcing.

This approach would again require localised community engagement and support, and as long as the values of the river warranted gull control, should be more socially acceptable. It is likely that there would be less opposition to local projects to protect specific values than wider regional control.

Individual values based control operations would require additional funding to current (or planned) mammalian predator control projects. Control operation cost would vary between rivers depending on the number of gulls. Depending on the size of the Southern Black-backed Gull/ Karoro population on each river section tentative costs are \$50,000 - \$220,000 for each river in Canterbury. As each project would be working independently, it is likely (especially in projects on lower river reaches) that ongoing immigration from surrounding rivers would require ongoing maintenance.

6.2 Control coordination

Currently there is a number of Southern Black-backed Gull/ Karoro control operations occurring throughout Canterbury, from the Mackenzie Basin to the Clarence River Mouth.

However, as discussed, current Southern Black-backed Gull/ Karoro control operations are currently being undertaken with no coordination. In order to deliver a regional Southern Black-backed Gull/ Karoro strategy there is a clear need for a Regional Coordinator.

A present there is no central depository for information and advice on undertaking Southern Black-backed Gull/ Karoro control, and each project is essentially learning as they go. During meetings with agencies undertaking Southern Black-backed Gull/ Karoro control there was a clear message that a centralised system with regional coordination was the best way forward.

The role of a Regional Coordinator would be to provide specialist advice;

- Provide advice during operation planning. This should include site visits to determine the best control method to utilise.
- Provide advice and material to assist the advocacy campaign around control operations to assist public engagement.
- Maintain and update SOP to latest learnings.
- Guide research needs and report findings to control practitioners.

6.3 Technical control method plan

Throughout Canterbury there is a number of control operations being carried out which are utilising a range of control methods. Alpha Chloralose and shooting are the main methods used, with some egg sterilisation also occurring (often in conjunction with other control methods).

Throughout discussions with control operators throughout Canterbury, there was a clear message that together with a regional strategy there is need for the development of best practise guidelines for Southern Black-backed Gull/ Karoro control.

There is a need for a review of control methods, both in relation to operational procedures, but also effectiveness. Then the development of guidelines and standard operating processes for control operations.

In particular there is a need to review the use of Alpha Chloralose, and especially the requirements for its use. Alpha Chloralose is the most commonly used method to Southern Black-backed Gull/ Karoro, but there is a number of differences and inconsistencies in its use throughout Canterbury. In

particular there was a desire for the application process and requirements to be streamlined to help with the planning and application process.

There was widespread agreement that there is a need for a more detailed technical control methods strategy which complements and would then help deliver a Canterbury Southern Black-backed Gull/ Karoro Strategy, and we strongly recommend the development of such a Technical Control Method Plan.

6.4 Research

The development of a regional strategy has been hampered by the lack of knowledge in several key areas. As such making informed decisions on the best approach moving forward has been challenging. Filling these information gaps is a key aspect of this strategy, and we identify several significant research requirements.

6.4.1 Southern Black-backed Gull/ Karoro foraging behaviour

Little is known about the daily foraging behaviour and range of Southern Black-backed Gull/ Karoro. With no in-depth studies undertaken in New Zealand, foraging range has just been estimated up to 50km, but there has been no detailed studies to investigate this.

Likewise, the foraging behaviour of gulls breeding on braided rivers have not been studied. With no knowledge on how gulls utilise both the river, and surrounding environments (farmland or urban centres). As such there is no measure how this relates to the impacts on braided river dependent shorebirds.

Research into the foraging behaviour and range of Southern Black-backed Gull/ Karoro breeding on braided rivers is recommended. A GPS tracking study Southern Black-backed Gull/ Karoro during the shorebird breeding season should be undertaken to determine the foraging regime of gulls. In particular this study should aim to investigate how Southern Black-backed Gull/ Karoro are using the braided river and other adjacent habitats.

6.4.2 Effectiveness of control methods

A range of control methods have been employed to reduce Southern Black-backed Gull/ Karoro numbers at colonies. At present the majority of these control operations have worked independently from each other.

In order to assess the effectiveness (including cost effectiveness) a research component should be incorporated into all control programmes. Ensure standardised recording so that different operations can be compared and evaluated.

Results of operations and learnings should be made available to others, and a forum established to link such practitioners (see above section relating to control coordination).

6.4.3 Immigration rates and direction post control

Southern Black-backed Gull are known to disperse some distance from natal colonies (Rowe 2013). However, despite multiple Southern Black-backed Gull/ Karoro control operations occurring in Canterbury, there has been no study looking at immigrating post control.

With significant Southern Black-backed Gull/ Karoro populations on many Canterbury rivers, there is no knowledge of the effects of removing colonies, and the immigration rate of birds into these colonies. Does removal of breeding birds create a “sink” where immigrating birds immediately fill vacated territories?

Inland rivers have fewer Southern Black-backed Gull/ Karoro, and one recommendation is to potentially treat these as containment areas. However immigration rates may need to be determined to better understand how to maintain these areas gull free.

Research into the immigration rates of Southern Black-backed Gull/ Karoro into control areas is recommended. To achieve this would require a large scale banding project in colonies nearby current or planned control operations. Again, this would require coordination at a regional scale, (see above section relating to coordination)

6.5 Public perception

Although there is a growing body of evidence which highlights the negative impacts Southern Black-backed Gull/ Karoro on braided river dependent shorebirds, this is not well known by the wider public.

There needs to be greater emphasis by braided river managers to ensure that the results of their work are disseminated to the wider public – not just the conservation movement. In our modern world there is numerous platforms that can be utilised for this. The aim should be to ensure that we tap into mainstream media to be sure a wide reach is achieved.

In addition to the impacts of Southern Black-backed Gull/ Karoro, further work to educate the public on the reasons for their control is warranted. Currently all control operations have some level of resistance from the public, either concerns over the need to control gulls, or the method employed (often an issue of toxin use). As more gull control is undertaken, it is likely that these issues are only going to increase. A specific marketing campaign to influence public perception on the conservation concerns and need to control Southern Black-backed Gull/ Karoro is recommended.

6.6 Land use

Southern Black-backed Gull/ Karoro numbers in Canterbury are not evenly distributed, with significantly greater numbers in low lying coastal areas. This distribution is almost entire driven by high intensity agriculture throughout this area. This land use has provided gulls with an abundant food source and enable their population numbers to explode. Furthermore poor waste management practises at landfills have added to this, helping to drive population growth.

Land use needs to be taken into consideration when developing a Canterbury wide strategy. With Southern Black-backed Gull/ Karoro having high breeding success and population growth in low lying areas, any control regime may be compromised. A significant part of public opposition to gull control is that gulls have managed to adapt well to human induced change, and that the problem is entirely human caused. There is increasing calls for better land and waste management that will readdress these issue into the future, so the need for lethal control options are not required in the future.

Although intensive agriculture will remain as the primary land use in all low lying areas, efforts should be made to develop ways that this land use will no longer be able to support such high populations and even enable long term reductions in gull numbers. Pivotal to this will be ensuring that waste management practises are improved, especially at landfills. Through their regulatory powers Environment Canterbury should ensure that all future resource consents take into consideration the potential food sources for Southern Black-backed Gull/ Karoro and ways to minimise this.

Land use change in upper river reaches is potentially the single biggest threat to Canterbury's shorebird populations. The move to intensive agricultural in the high country is likely to see an improvement in breeding success for Southern Black-backed Gull/ Karoro, and lead to significant population increases. Left unchecked this will almost certainly cause declines in threatened shorebird populations.

The current expansion of irrigation and dairy farming in the Mackenzie Basin highlights the potential risks to braided river dependent shorebirds. Increased irrigation and stock numbers will provide improved foraging for Southern Black-backed Gull/ Karoro, and will almost certainly see an increase in their numbers. Walker and Monks (2017) concluded that intensive land use was one of the key drivers causing endemic shorebirds to retreat into the higher altitude rivers.

Environment Canterbury should not encourage agriculturally intensification in the high country, and through its regulatory powers make the relevant plan changes to ensure that increased irrigation and land intensification are not permissible.

6.7 Tangata Whenua – Mahinga Kai

Southern Black-backed Gull/ Karoro eggs and chicks have been a traditional food of Ngāi Tahu, and there is a number of Hapu that have expressed interest in resuming the collection of eggs and chicks for a food source.

Under current legislation there is nothing preventing the collection of eggs or chicks, as Southern Black-backed Gull/ Karoro are an unprotected species.

The removal of eggs and chicks as Mana Kai is unlikely to provide a significant reduction in population numbers. However, it could form a part of a long term management programme aimed at reducing population growth, and/or immigration into control areas. The targeted collecting of eggs and chicks from specific colonies, or regions could help reduce overall breeding success and limit population growth.

The resumption of use of Southern Black-backed Gull/ Karoro eggs for Mana Kai is recommended, and should be encouraged. In order to promote this, greater communication of the location and size of Southern Black-backed Gull/ Karoro colonies should occur between Environment Canterbury and Tangata Whenua. Potentially the Te Paiherenga forum is the best platform for this to occur.

7. Acknowledgements

This project would not have been possible without the input of a huge range of people. In particular Frances Schmechel from Environment Canterbury who has guided this process and was invaluable throughout. Donna Lil provided support with community engagement and facilitated public meetings, which was of immense help to this project. During this project a wide range of people provided input, and we thank everybody who sent us their thoughts and/or attended meetings. Although too many to name individually a number of people contributed significantly. We are grateful for the significant contributions from Andrew Crossland, Neal Dean, Simone Cleland, Courtney Popenhagen, Donna Field, Manaia Cunningham, Richard Maloney, Andy Grant and Sonny Whitelaw. Thanks to Patrick Crowe and Rachel Wilson (both WMIL) for producing the maps and other data analysis.

8. References

- AECOM. 2016. Permanent disposal of earthquake waste at the Burwood Recourse Recovery Park (Site B) – Water Quality Report. Retrieved from <http://files.ecan.govt.nz/public/consent-projects/burwood/Appendix B - Water Quality Report.PDF>
- Avisure. 2016. Christchurch International Airport Ltd Off-Airport Bird Hazard Management Plan. Report to CIAL.
- Belant, J.L. 1997. Gulls in urban environments: landscape-level management to reduce conflict. *Landscape and urban planning* 38: 245-258.
- Bell, M.D. 2018. Waiau Toa/ Clarence River mouth black-billed gull outcome monitoring 2016-2018. Unpublished Wildlife Management International Technical Report to the Department of Conservation and Environment Canterbury.
- Bell, M., McArthur, N., and Harborne, P. 2018. Clarence River black-fronted tern restoration project – 2017/2018 operational report. Client report prepared for the Department of Conservation. Wildlife Management International Ltd, Blenheim.
- Bell, M., and Harborne, P in prep. 2018. The identification of predators at black-billed gull (*Larus bulleri*) colonies using time lapse photography.
- Biswell, S.F. 2005. Southern black-backed gulls. *New Zealand Geographic* 73: pp 46-61.
- Brown, K.; Keedwell, R. 1998. Alphachloralose control operations at southern black backed gull colonies in the Tekapo and Ahuriri Valleys. Canterbury Conservancy Technical Report Series Number 7, Department of Conservation, Twizel.
- Clark, G.; Fenwick, S.; Boxall, N.; Swanney, S.; Nicol, C. 1999: *Salmonella* Brandenburg abortions in sheep, pathogenesis and pathology. *Society of sheep & beef cattle veterinarians, New Zealand veterinary association annual seminar* 29: 12-22.
- Clark, G. 2000. Spread of *Salmonella* Brandenburg organisms in sheep yards. MAF Policy project report.
- Clark, R.G. 2001. Black-backed gulls (*Larus dominicanus*) and their role in the spread of *Salmonella* Brandenburg. *Proceedings of the New Zealand Society of Animal Production* 61: 71-73
- Corkhill, P. (1970) Effect of egg puncturing on limiting chick production in Herring Gulls. *Bird Study* 17: 340-342.
- Fenlon, D.R. 1981: Seagulls (*Larus spp.*) as vectors of salmonellae: an investigation into the range of serotypes and numbers of salmonellae in gull faeces. *Journal of hygiene (Cambridge)* 86: 195-202.
- Fordham, R.A. 1964. Breeding biology of the southern black-backed gull. II: Incubation and the chick stage. *Notornis* 11: 110–126.

- Fordham, R. A. 1967. History and status of the Dominican gull in Wellington. *Notornis* 14: 144-153.
- Fordham, R.A. 1968. Dispersion and dispersal of the Dominican gull in Wellington, New Zealand. *New Zealand Ecological Society* 15: 40-50.
- Fordham, R.A. 1985. Dominican gull. In: *Readers Digest complete book of New Zealand birds*. Sydney: Readers Digest. 231 p.
- Hatch, J.J., 1996. Threats to public health from gulls (Laridae). *J. Environ. Health Res.* 6, 5-16.
- Heather, B.D. & Robertson, H.A. 1996. *Field guide to the birds of New Zealand*. Auckland: Viking. 432 p.
- Heather, B.D. & Robertson, H.A. 2005. *The field guide to the birds of New Zealand*. Penguin Books, Auckland. pp 122, 123, 332-334.
- Higgins, P.J.; Davies, S.J.J.F. (Eds). 1996. *Handbook of Australian, New Zealand and Antarctic birds. Volume 3: Snipe to pigeons*. Melbourne: Oxford University Press.
- Horton, N., Brough, T.; Rochard, J.B.A. 1983. The importance of refuse tips to gulls wintering in an inland area of south-east England. *Journal of Applied Ecology* 20: 751-765.
- Galbraith, M. 2015. Changes in the breeding status of the southern black-backed gull (*Larus dominicanus*) colonies on Rangitoto Island, Hauraki Gulf, New Zealand. *Notornis* 62: 192-201.
- Hutching, G. 2006. Gulls, terns and skuas. Te Ara – the Encyclopedia of New Zealand. <http://www.TeAra.govt.nz/en/photograph/6124/controlling-gulls-near-airports> (accessed 26 June 2018)
- Johnston, W.S.; MacLachlan, G.K.; Hopkins, G.F. 1979: The possible involvement of seagulls (*Larus* sp) in the transmission of Salmonella in dairy cattle. *Veterinary record* 105: 526-527.
- Jones, F., Smith, P., Watson, D.C., 1978. Pollution of a water supply catchment by breeding gulls and the potential environmental health implications. *J. Inst. Water Eng. Sci.* 32, 469- 482.
- Maloney, R. F. 1999. Bird populations in nine braided rivers of the Upper Waitaki Basin, South Island, New Zealand: changes after 30 years. *Notornis* 46: 243-256.
- McDiarmid, A. 1962. Diseases of free-living wild animals. *Agric. Stud* 57: 8-28, Food & Agriculture Organization, Rome.
- Miers, K.H and Williams, M., 1969. Nesting of the black swan at Lake Ellesmere, New Zealand. *Wildfowl*, 20(20), pp. 23-32.
- Mischler, C.P.; Bell, M.D. 2016. Waiiau Toa/ Clarence River Black-billed Gull Management. Unpublished technical report by Wildlife Management International Ltd to Environment Canterbury.
- Miskelly, C.M. 2013. Southern black-backed gull. In: Miskelly, C.M. (ed.) *New Zealand Birds Online*: www.nzbirdsonline.org.nz
- Monaghan, P., Shedden, C.B., Fricker, CR., Girdwood, R.W.A., 1985. **Salmonella** carriage by herring gulls in the Clyde area of Scotland in relation to their feeding ecology. *J. Appl. Ecol.* 22, 669-680.
- Moorhouse, R. 2017. Results of Nelson Nature, Environmental Monitoring – Coastal Habitats – Shorebird Survey – October 2016 to January 2017.
- More, E.; Ayats, T.; Ryan, P.G.; Naicker, P.R.; Keddy, K.H.; Gaglio, D.; Witteveen, M.; Cerda-Cuellar, M. 2007. Seabirds (*Laridae*) as a source of *Campylobacter* spp., *Salmonella* spp. and antimicrobial resistance in South Africa. *Environmental Microbiology* 19 (10): 4164 – 4176
- Nelson, P. C. 1994. Bird control in New Zealand using Alpha-chloralose and DRC-1339. *Proceedings of the Sixteenth Vertebrate Pest Conference* 40: 259 - 264
- Nelson, M., Jones, S. H., Edwards, J.C., Ellis, C. 2008. Characterization of *Escherichia coli* populations from gulls, landfill trash, and wastewater using ribotyping. *Diseases of Aquatic Organisms* 81: 53-63.

- Nicoll, J. 2017. Porirua tip-goer claims gull left 'like wounded soldier' after being shot. Retrieved from <https://www.stuff.co.nz/environment/93361252/porirua-tipgoer-claims-gull-left-like-wounded-soldier-after-being-shot>
- O'Donnell, C.F.J.; Sanders, M.; Woolmore, C.; Maloney, R.F. 2016. Management and research priorities for conserving biodiversity on New Zealand's braided rivers. Department of Conservation, Wellington New Zealand, 46p.
- Oliver, W.R.B. 1955. *New Zealand birds*. Wellington: A.H. & A.W. Reed.
- Oliver, R. 1973. *Studies of the breeding biology of the southern black-backed gull on Rangitoto Island*. MSc thesis, University of Auckland, New Zealand.
- O'Neil, A. 2013. Seagulls threatening businesses <http://www.stuff.co.nz/dominion-post/news/local-papers/kapi-mana-news/9254759/Seagulls-threatening-businesses?rm=m>
- Powlesland, R.G.; Robertson, H.A. 1987. Changes in gull numbers over 25 years and notes on other birds of the Otaki-Ohau coast. *Notornis* 34: 327-338.
- Quessey, S., Messier, S. 1992. Prevalence of *Salmonella* spp., *Campylobacter* spp. and *Listeria* spp. in ring-billed gulls (*Larus delawarensis*). *J. Wildl. Dis.* 28, 526 - 531.
- Raven, S.J and Coulson, J.C. 1997. The distribution and abundance of *Larus* gulls nesting on buildings in Britain and Ireland. *Bird Study*, 44: pp 13-34.
- Reed, K.D., Meece, J.K., and Henkel, J.S. (2003) Birds, Migration and Emerging Zoonoses: West Nile Virus, Lyme Disease, Influenza A and Enteropathogens. *Clin Med Res* 1: 5–12.
- Reilly, W.G., Forbes, G.I., Paterson, G.M., Sharp, J.C.M., 1981. Human and animal salmonellosis in Scotland associated with environmental contamination, 1973-1979. *Vet. Rec.* 108, 553-555.
- Robertson, C.J.R. 1964a. 14th annual report of the New Zealand bird banding scheme for the year ended 31st March 1964. Dominion Museum, Wellington. P24.
- Robertson, C.J.R. 1964b. Observations on black-backed gull predation at the Cape Kidnappers gannetries 1959-1963. *Notornis* 10: 393-403.
- Robertson, C.J.R. 1972. Preliminary report on bird banding in New Zealand 1964-1971. *Notornis* 19: 61-73.
- Robertson, C.J.R. 1973. Preliminary report on bird banding in New Zealand 1971-1972. *Notornis* 20: 59-70.
- Robertson, C.J.R.; Hyvönen, P.; Fraser, M.J.; Pickard, C.R. 2007. *The atlas of bird distribution in New Zealand 1999-2004*. Wellington: Ornithological Society of New Zealand Inc.
- Robertson, H.A. 1992. Trends in the numbers and distribution of coastal birds in Wellington Harbour. *Notornis* 39(4): 263-289.
- Rowe, K.R. 2013 Dispersal of southern black-backed gulls (*Larus dominicanus dominicanus*) banded in Canterbury, New Zealand, 1959-1993. *Notornis* 60: 134-142
- Schlesselmann, A.V., O'Donnell, C.F.J., Monks, J.M., Robertson, B.C. 2018. Clearing islands as refugia for black-fronted tern (*Chlidonias albostratus*) breeding colonies in braided rivers. *New Zealand Journal of Ecology*
- Solman, V.E.F. 1994. Gulls. *The Handbook: Prevention and control of wildlife damage*. 62.
- Steffens K.E, Sanders M.D, Gleeson D.M, Pullen K.M, Stowe C.J 2012. Identification of predators at black-fronted tern *Chlidonias albostratus* nests, using mtDNA analysis and digital video recorders. *New Zealand Journal of Ecology* 36: 48–55.
- Thierry, A., Dutton, P. and Popenhagen, C. 2016. Waimakariri River Regional Park Black-billed Gull Management 2015-2016 Breeding Season. Unpublished report, Keystone Ecology and Environment Canterbury Report.
- Turbott, E.G. 1969. Roof nesting of black-backed gulls. *Notornis* 16: 187-9.

- USDA. 2001. DRC-1339 (Starlicide). USDA National Wildlife Research Centre – Staff Publications. 740. Retrieved from http://digitalcommons.unl.edu/cgi/viewcontent.cgi?article=1735&context=icwdm_usdanwrc
- Williams, M.J. 1963. Predatory habits of the black-backed gull. *Notornis* 10: 239-240.
- Williams, M., Gummer, H., Powlesland, R., Robertson, H., and Taylor, G. 2006. Migration and movements of birds to New Zealand and surrounding seas. Department of Conservation, Wellington, New Zealand.
- Wills, D.E.; Murray, J.; Powlesland, R.G. 2003. Impact of management on the breeding success of the northern New Zealand dotterel (*Charadrius obscurus aquilonius*) on Matakana Island, Bay of Plenty. *Notornis* 50: 1-10.
- Winfiel, M.D., and Groisman, E.A. (2003) Role of Nonhost Environments in the Lifestyles of Salmonella and E. coli. *App Environ Microbial* 69: 3687–3694.
- Yorio, P and Quintana F. 1997. Predation by Kelp Gulls *Larus dominicanus* at a mixed-species colony of Royal Tern *Sterna maxima* and Cayenne Terns *eurygnatha* in Patagonia. *IBIS* 139: 536-541.
- Yorio, P.; Bertellotti, M.; Gandini, P.; Frere, E. 1998. Kelp gulls *Larus Dominicanus* breeding on the argentine coast: population status and relationship with coastal management and conservation. *Marine Ornithology* 26: 11-18.